

# Galaxy-Scale AGN Outflows: Two Puzzles, Two Solutions

Claude-André Faucher-Giguère


UC Berkeley

Miller Institute for Basic Research in Science

with Eliot Quataert & Norm Murray

# Observational breakthroughs on AGN outflows

- Massive, galaxy-scale AGN outflows in local ULIRGs
  - ➔ neutral, ionized, CO, OH, HCN, ...
- Herschel, E-VLA, ALMA, ... to revolutionize this field
- Physical conditions in QSO outflows using low-ion BALs ( $\Rightarrow$  energetics)



- Photoionization modeling particularly constraining in QSOs with FeII\* broad line absorption ( $T \sim 10^4$  K,  $v \sim 5,000$  km/s; FeLoBALs):

## Compact absorbers must form *in situ*, at $R \sim \text{kpc}$ from SMBHs

- If they traveled from the SMBH to their implied location...

$$t_{\text{flow}} \approx \frac{R}{v} \approx 3 \times 10^5 \text{ yr} \left( \frac{R}{3 \text{ kpc}} \right) \left( \frac{v}{10,000 \text{ km s}^{-1}} \right)^{-1}$$


- But destroyed by hydro instabilities and thermal evaporation in

$$t_{\text{KH}}, t_{\text{evap}} \sim \text{few} \times 10^3 \text{ yr}$$

**Not a direct accretion disk wind!**

## Radiative shock model


- Form in interaction of the QSO blast wave with an ISM clump:



pre-existing ISM cloud

## Cloud crushing by shocks, Kelvin-Helmholtz instability

- Well-studied problem for SNRs (e.g., Klein+94, Cooper+09)



## Requirements for radiative shocks explain properties of cool absorbers

- Acceleration, cold gas:

$$\frac{t_{\text{drag}}}{t_{\text{cool}}} < \frac{t_{\text{KH}}}{t_{\text{cc}}} \Rightarrow N_{\text{H}} \gtrsim 10^{20} \text{ cm}^{-2} \left( \frac{v_{\text{sh}}}{5,000 \text{ km s}^{-1}} \right)^{4.2}$$

- Post-shock compression:

$$n_{\text{H}}^{\text{BAL}} \approx 4n_{\text{H}}^{\text{pre}} \left( \frac{T_{\text{sh}}}{10^4 \text{ K}} \right) \sim 10^4 \text{ cm}^{-3}$$

$$\Rightarrow \Delta R \sim N_{\text{H}}/n_{\text{H}} \sim 0.01 \text{ pc}$$

- Also: super-thermal line widths, multiple v components, reddening, ...

# Energetics of QSO outflows

- Outflows are multiphase
- Most of kinetic power in hot flow:


$$\dot{M}_{\text{hot}} = 8\pi\Omega_{\text{hot}} R N_{\text{H}}^{\text{hot}} \mu m_{\text{p}} v_{\text{hot}}$$

- Using radiative shock model:

$$\dot{E}_{\text{k}} \approx 2 - 5\% L_{\text{AGN}}$$


$$\dot{M} \approx 1,000 - 2,000 M_{\odot}/\text{yr}$$

$$\dot{P} \approx 2 - 10 L_{\text{AGN}}/c$$





## The puzzle of large momentum fluxes



- If all photons scatter once &  $P$  is conserved,

$$\dot{P} \sim L_{\text{AGN}}/c$$

- Observations indicate

$$\dot{P} \sim 10L_{\text{AGN}}/c$$

- Simulations also require

$$\dot{P} \gg L_{\text{AGN}}/c$$

to reproduce  $M_{\bullet}-\sigma$  (DeBuhr+)

## Momentum driving

$$t_{cool} \ll t_{flow}$$

No thermal pressure

$$P_{final} \sim P_{start}$$

e.g., AGB wind


## Energy driving

$$t_{cool} \gg t_{flow}$$

Shocked gas does work


$$P_{final} \gg P_{start}$$

e.g., Sedov-Taylor SNR



## Proposal: AGN outflows are energy-driven

- Possible in ULIRGs despite extreme densities
  - ➔ relevant criterion is cooling of reverse shock:  $T_{sw} \sim 10^{10}$  K for  $v_{in} \sim 0.1c$
  - ➔ 2-T plasma inhibits IC cooling



CAFG & Quataert, in prep.


# Energy conservation naturally explains measured AGN momentum boosts

- Predicts

$$\frac{\dot{P}}{L_{\text{AGN}}/c} \sim \frac{1}{2} \left( \frac{\text{nuclear wind speed}}{\text{galaxy wind speed}} \right)$$

- To be tested soon with Herschel, E-VLA, ALMA, ...

- Analytic model will inform numerical implementations



## Summary

- Compact, cool absorbers form in radiative shocks
- Energetics in good agreement with  $M_{\bullet}$ - $\sigma$  requirements
- Observations of galaxy-scale AGN outflows suggest

$$\dot{P} \gg L_{\text{AGN}}/c$$

- Proposal: outflows are energy-conserving
- Prediction:

$$\frac{\dot{P}}{L_{\text{AGN}}/c} \sim \frac{1}{2} \left( \frac{\text{nuclear wind speed}}{\text{galaxy wind speed}} \right)$$



## Robust to mixing, leakage


- Stellar wind bubbles smaller & slower than in energy-conserving models (Castor)
  - ➡ cooling due to mixing (McKee+84)
  - ➡ hot gas vents out (H.-C. & Murray 09)
- AGN winds more robust
  - ➡  $\sim 30\times$  wind mass of cool gas before catastrophic ff cooling
  - ➡ escape along paths  $< 10^{-3}$  underdense can still boost  $P$  by factor  $> 10$  in ULIRGs

Carina nebula




# The possible roles of AGN feedback

Establish correlations between  
SMBH and galaxy properties



Truncate star formation



Models that assume  $f \sim 5\%$   $L_{\text{AGN}}$  couples to ISM are successful in  
explaining in these observations