

Astro2020 Science White Paper

Closing Gaps in Our Astrochemical Heritage: From Molecular Clouds to Planets

Thematic Areas: Planetary Systems Star and Planet Formation
 Formation and Evolution of Compact Objects Cosmology and Fundamental Physics
 Stars and Stellar Evolution Resolved Stellar Populations and their Environments
 Galaxy Evolution Multi-Messenger Astronomy and Astrophysics

Principal Author:

Name: Brett A. McGuire

Institution: National Radio Astronomy Observatory

Email: bmcguire@nrao.edu

Phone: (434) 244-6807

Co-authors: Edwin Bergin (University of Michigan), Geoffrey A. Blake (California Institute of Technology), Andrew M. Burkhardt (Harvard-Smithsonian Center for Astrophysics), P. Brandon Carroll (Harvard-Smithsonian Center for Astrophysics), L. Ilse-dore Cleeves (University of Virginia), Quentin Kral (Paris Observatory), Ryan A. Loomis (National Radio Astronomy Observatory), Klaus M. Pontoppidan (Space Telescope Science Institute), Christopher N. Shingledecker (Max Planck Institute for Extraterrestrial Physics), Eric. R. Willis (University of Virginia)

Abstract (optional): One of the primary mechanisms for inferring the dynamical history of planets in our Solar System and in exoplanetary systems is through observation of elemental ratios (i.e. C/O). The ability to effectively use these observations relies critically on a robust understanding of the chemistry and evolutionary history of the observed abundances. Significant efforts have been devoted to this area from within astrochemistry circles, and these efforts should be supported going forward by the larger exoplanetary science community. In addition, the construction of a next-generation radio interferometer will be required to test many of these predictive models *in situ*.

1 Background

The location of condensation fronts (snow lines) in protoplanetary disks (PPDs) can have substantial effects on the physical and chemical evolution of planets in the system. Indeed, many models predict that these condensation fronts are the locations where planetesimal formation begins (Ros & Johansen 2013; Schoonenberg & Ormel 2017; Drazkowska & Alibert 2017). The location of these condensation fronts is determined both by the incident radiation from the protostar, but also, in large part, by the chemical composition of the gas and ice. The desorption temperature of the bulk ice varies significantly with the chemical composition and mixing ratio (Collings et al. 2004), even of minor constituents. This can be further complicated by a variance in the sticking efficiency of the icy grains with composition (Wang et al. 2005). It is clear that the underlying chemical composition in these ices, even at low mixing ratios, can have a substantial effect on the efficiency of planet formation.

Further, the relative gas vs solid-phase inventories at snow lines govern the composition of the planets and their atmospheres. These compositions (e.g., the C/O ratio in planetary atmospheres; Öberg et al. 2011) may therefore be useful probes of the history of exoplanets. Effectively using observed exoplanet elemental ratios to test those models therefore relies heavily on our ability to accurately interpret chemical abundances measured in disks to infer the inventory of undetectable species, and how these measurements affect observed elemental ratios.

Constraining the chemical compositions in PPDs is challenging. The most complex gas-phase species yet detected in PPDs are methanol (CH_3OH ; Walsh et al. 2016) and methyl cyanide (CH_3CN ; Öberg et al. 2015), and more complex molecules are unlikely to be routinely observed (Walsh et al. 2014). Further, in cold or shielded regions within disks, ices may be the major repository of C and O (Bergin et al. 2016). Direct observation of ices typically require space-based infrared facilities and a target source with a sufficiently bright background continuum *and* favorable viewing geometry. Direct observations of bulk water ice in any disk, in emission, is possible using the strong $43\ \mu\text{m}$ ice band, and to a lesser degree, the weaker $62\ \mu\text{m}$ ice band (McClure et al. 2015; Kamp et al. 2018). SOFIA-HIRMES will access these bands for the brightest disks (Richards et al. 2018), but to obtain a census of the water ice mass in disks around lower-mass stars, a sensitive far-infrared space telescope is required, such as ESA's SPICA concept, or the Origins Space Telescope.

Even in massive young stellar objects with substantial ice columns, the list of firmly-detected ice species numbers less than a dozen (Boogert et al. 2015). Spitzer provided basic constraints on the typical chemistry of bulk ice species in particular in low-mass star-forming regions, with the largest detected ice species being CH_3OH (Boogert et al. 2015). JWST will build on this to obtain measurements of more complex organic species both in the quiescent interstellar medium, and in protostars (e.g., in the IceAge Early Release Science program, McClure et al. 2017). Casting an even wider net, SPHEREx will observe ices in roughly 100 times as many sources, including molecular clouds, young stellar objects, and protoplanetary disks, as have been observed to date. This will allow for the quantification of the principal ice components across an enormous sample of sources. Our knowledge of gas and ice compositions in disks, however, must rely on astrochemical models, using the detectable simple gas species as the primary constraint.

These models are not yet robust, despite substantial efforts (Garrod 2013; Herbst 2017; Wakelam et al. 2015). To reliably infer chemical compositions in PPDs, in which chemistry is closely coupled to physical conditions (temperature, density, UV flux, evolving dust populations, etc.) that vary with radius and scale height, necessitates that these models be well-constrained by observations in earlier stages of star formation, where more robust inventories can be observed. *Here, we briefly discuss the state-of-the-art in earlier stages of chemical evolution, and outline future directions and the technology and community support needed to enable them.*

2 Chemical evolution in quiescent environments

Cold ($T_{ex} \sim 10$ K), dark, starless clouds are an ideal environment in which to test the time evolution of chemical reaction networks under ‘constant’ physical conditions. The typical sources for exploring this stage of star and planet formation are highly extended ($\theta_s > 30''$). At these excitation temperatures, most molecules, especially those more complex than CH_3OH , will display bright emission at cm wavelengths. The best strategy for obtaining the required abundance and excitation information is therefore deep, broadband spectral line surveys at cm-wavelengths using single-dish facilities combined with chemical modeling (Cordiner et al. 2017; Kaifu et al. 2004; McGuire et al. 2018). The best extant instrument for this work is the Green Bank Telescope (GBT).

Even in a source as extended as TMC-1, there exists evidence for large chemical variation over small spatial scales (0.02–0.03 pc; Peng et al. 1998; Dickens et al. 2001), and observations of such would probe the effects of the very earliest stages of collapse on the chemistry. These observations will require high spatial resolution, high surface-brightness sensitivity, and broad spectral bandwidth to be efficient. Because of the cold temperatures, this further requires a cm-wavelength facility. While the VLA satisfies this need, it lacks the combination of surface brightness sensitivity and bandwidth necessary to conduct a sensitive line survey with any efficiency. The proposed ngVLA, in its current conceptual design, would prove more than optimal for this task.

3 ... and in complex, star-forming regions

Sgr B2(N), Orion-KL, and NGC 6334I are examples of the most chemically and physically complex stages of star formation observed and modeled (Garrod et al. 2008; Crockett et al. 2014; McGuire et al. 2017). The extreme physical environments, interplay between the gas and ice, and complex structures often result in models which can reproduce observed abundances in, at best, a qualitative fashion (Laas et al. 2011; Garrod et al. 2008; Garrod 2013). These types of physical and chemical interactions are the same that are at play in PPDs, albeit on a much larger scale. It is therefore these environments that provide a unique test environment for the similar PDR style models applied to PPDs. Improvements in these models are most likely to be achieved by testing them against a wide range of environments from quiescent clouds to massive star forming regions.

Existing facilities in the (sub-)millimeter regime, particularly ALMA, are capable of reaching the line-confusion limit in broadband line surveys of complex, star forming regions in a matter of hours. That is, the density of molecular lines in the resulting spectra is such that there is at least one spectral line every FWHM; there are no line-free channels. When line-confusion is reached, no

additional information can be gained from longer integration times at the same spatial resolution. We are rapidly approaching the point at which deep drill observations with ALMA and the GBT will no longer produce new spectral lines. The end result of this limit is that no further detections of new species can be made, and thus the refinement of chemical networks will be limited to those species detectable at the confusion limit.

In the cm-wave regime, however, the most sensitive survey, the Prebiotic Interstellar Molecular Survey (PRIMOS; Neill et al. 2012) with the most sensitive facility (the GBT) toward the most line-rich source (Sgr B2) has lines in only $\sim 10\text{-}20\%$ of its channels at cm-wavelengths. Diminishing returns on sensitivity for this survey have already set in, and to increase the sensitivity substantially would require hundreds of hours of dedicated observations for every ~ 2 GHz window. Further, ALMA observations at higher frequencies are revealing that the region displays widely variable physical conditions within a relatively small environment, variation that is lost to single-dish GBT observations (Belloche et al. 2016). This offers a valuable opportunity to probe environment-induced chemical differentiation using spatially resolved observations which are not possible with the GBT.

While the brightest lines in Sgr B2 are readily detectable with extant facilities such as the VLA and ATCA (Corby et al. 2015), the complex species are not efficiently observable, especially at high angular resolution. The only viable solution to this problem is the construction of a next-generation interferometer like the next-generation Very Large Array (ngVLA). Under the current design concept (Selina et al. 2018), the ngVLA is likely to not only be capable of spatially resolving chemistry in Sgr B2(N), but to reach the line-confusion limit even at cm-wavelengths, providing substantial new, complex molecular detections and their correlation with the physical conditions at work.

4 ... in protostellar environments

The protostellar stage of star formation spans a wide range of sources, from ‘hot corinos’ such as IRAS 16293 (Jørgensen et al. 2016) to chemically rich outflow sources like L1157 (Mendoza et al. 2014). The molecular outflows from active young protostars like L1157 provide a unique environment to test the immediate impact of rapid changes in temperature and density on ‘pristine’ molecular clouds. In these sources, distance from the protostar along the outflow corresponds directly to time, allowing modeling of temporal evolution. Hot corinos are the next stage after L1157 and before hot cores like Sgr B2(N).

The best method for constraining the chemistry in these sources is broadband spectral line surveys. These sources are relatively compact, are beam-diluted in single-dish observations, and their linewidths are typically such that high-sensitivity data are often at or approaching line confusion limits, necessitating interferometric observation (Jørgensen et al. 2016). Because they are typically moderately warm (50–150 K; Jørgensen et al. 2016; McGuire et al. 2015), complex molecules display bright features in the 1–3 mm window well-covered by ALMA, NOEMA, and the SMA.

The SOLIS project at NOEMA and IRAM, which examines a sample of solar-type protostellar sources (Ceccarelli et al. 2017), and the PILS project at ALMA, which has focused particularly

on an exquisitely detailed examination of the IRAS 16293 system (Jørgensen et al. 2016), are the primary active dedicated investigations into this class of objects. Extant observational facilities are likely sufficient to provide the required insight. Ongoing and future efforts on ALMA, NOEMA, and the SMA examining a large sample of protostellar environments are critical to constraining the effects of physical environment on chemical evolution and incorporating these effects into models.

5 ... and in PPDs

A number of larger scale studies of disk physical properties and chemistry have been enabled by the sensitivity and resolution of ALMA. Efforts to measure fundamental properties such as bulk mass (Ansdell et al. 2016), and larger compositional studies examining nitrogen (Guzmán et al. 2017) and deuterium (Huang et al. 2017) ratios, are now possible for larger samples sizes. With robust chemical modeling, ALMA observations of simple hydrocarbons (e.g., C₂H) can perhaps be used to infer C/O ratios indirectly (Bergin et al. 2016; Cleeves et al. 2018).

JWST has the potential to directly estimate C/O ratios in inner disks (within 1 AU) through the rich ro-vibrational gas-phase spectra of simple organics and high-J rotational water lines known to be typical in disks around solar-mass stars (Pontoppidan et al. 2010; Carr & Najita 2011). For direct estimates of the total oxygen abundance in the regions around and beyond the water snowline (radii of 1-100 AU), a sensitive far-infrared space telescope, such as the Origins Space Telescope, is needed to measure the mass of bulk cold water ice and gas (Blevins et al. 2016). Access to the 112 μ m ground-state line of HD from will provide precise estimates of the total disk gas mass (Trapman et al. 2017), which is required as grounding information for any estimates of absolute abundances of rare species as observed by ALMA and potentially ngVLA. Looking even further along the evolutionary path to debris disks, ALMA observations are just now resolving the gaseous CO content released from exocometary collisions in systems such as β Pictoris (Kral et al. 2016; Matrà et al. 2016). While studies of molecular emission from ALMA are now possible, infrared observations with a mission such as Origins may provide the best access to probe this gaseous remnant of the molecular content, and critically the C/O ratio, during the epoch of planet formation.

Yet, these infrared facilities will still largely be blind to the complex chemistry occurring within the ices, and to the combined ice/gas coupling in warmer gas. That insight is only achievable through longer-wavelength radio observations of gas-phase species, that are sensitive to populations orders of magnitude lower in abundance, coupled with chemical modeling. Application of the insight gained from these models in the terrestrial planet forming region, near the H₂O snow line, will require AU-scale resolution with high sensitivity. A high-frequency ngVLA is the only facility currently in design with the required specifications. Further, while these models can then interpret results for C and O in disks, observations of the major N-bearing species, NH₃ will still be required at similar scales. An ngVLA operating at least through the 24 GHz NH₃ transition is therefore necessary. Further, such a facility must support broadband, high surface-brightness sensitivity capabilities at the small to modest (0.1–3'') scales required to probe the effects of the widely variable physical conditions on the chemical evolution in sources like Sgr B2 and unlock chemical inventories obscured by line confusion suffered at extant facilities like ALMA. The same instrument should be capable of observing NH₃ and imaging the planet forming zone inside the water snow line in disks, requiring AU-scale (mas) resolution capabilities.

Summary

The chemical evolution of all ranges of environments from planetary atmospheres ([Arsenovic et al. 2016](#)) to disk midplanes ([Cleeves et al. 2013](#)) to molecular clouds ([Hudson et al. 2008](#)) is impacted by radiation. The last few years have seen significant advances in modeling efforts to accurately include these effects ([Shingledecker & Herbst 2018](#)), but these efforts are young, and require substantial training on ‘simple’ environments like quiescent clouds where the radiation effects can be easily decoupled from dynamical physical events such as outflows and shocks. In short, efforts at the GBT and other large single-dish facilities are needed to survey quiescent chemical clouds. These efforts would be greatly bolstered by the development of broadband capabilities on the ngVLA to enable investigations of the chemical evolution upon initial gravitational collapse, and complemented with the infrared capabilities of JWST and the Origins Space Telescope.

A combination of observational and theoretical efforts are needed that are focused on placing rigorous constraints on models of chemical evolution at each stage of the star and planet formation process. The observational requirements of these investigations necessitate a commitment by the community to support extant facilities and the development of technological advancements for the same, as well as the development of a next-generation VLA with high sensitivity and exquisite spatial resolution. The ultimate goal should be producing models that can be used to better calibrate our estimates of gas and ice compositions in PPDs, given the observations of simple gas-phase species achievable in these sources, and the construction of a radio facility capable of observationally exploiting these models *in situ* during planet formation.

References

- Ansdell, M., Williams, J. P., van der Marel, N., et al. 2016, *The Astrophysical Journal*, 828, 1
- Arsenovic, P., Rozanov, E., Stenke, A., et al. 2016, *Journal of Atmospheric and Solar-Terrestrial Physics*, 149, 180
- Belloche, A., Müller, H. S. P., Garrod, R. T., & Menten, K. M. 2016, *Astronomy and Astrophysics*, 587, A91
- Bergin, E. A., Du, F., Cleeves, L. I., et al. 2016, *The Astrophysical Journal*, 831, 1
- Blevins, S. M., Pontoppidan, K. M., Banzatti, A., et al. 2016, *The Astrophysical Journal*, 818, 1
- Boogert, A. C. A., Gerakines, P. A., & Whittet, D. C. B. 2015, *Annual Review of Astronomy and Astrophysics*, 53, 541
- Carr, J. S., & Najita, J. R. 2011, *The Astrophysical Journal*, 733, 102
- Ceccarelli, C., Caselli, P., Fontani, F., et al. 2017, *The Astrophysical Journal*, 850, 176
- Cleeves, L. I., Adams, F. C., Bergin, E. A., & Visser, R. 2013, *The Astrophysical Journal*, 777, 28
- Cleeves, L. I., Öberg, K. I., Wilner, D. J., et al. 2018, *The Astrophysical Journal*, 865, 0
- Collings, M. P., Anderson, M. A., Chen, R., et al. 2004, *Monthly Notices of the Royal Astronomical Society*, 354, 1133
- Corby, J. F., Jones, P. A., Cunningham, M. R., et al. 2015, *Monthly Notices of the Royal Astronomical Society*, 452, 3969
- Cordiner, M. A., Charnley, S. B., Kisiel, Z., McGuire, B. A., & Kuan, Y.-J. 2017, *The Astrophysical Journal*, 850, 187
- Crockett, N. R., Bergin, E. A., Neill, J. L., et al. 2014, *The Astrophysical Journal*, 787, 112
- Dickens, J. E., Langer, W. D., & Velusamy, T. 2001, *The Astrophysical Journal*, 558, 693
- Drazkowska, J., & Alibert, Y. 2017, 608, A92
- Garrod, R. T. 2013, *The Astrophysical Journal*, 765, 60
- Garrod, R. T., Weaver, S. L. W., & Herbst, E. 2008, *Astrophys. J.*, 682, 283
- Guzmán, V. V., Öberg, K. I., Huang, J., Loomis, R., & Qi, C. 2017, *The Astrophysical Journal*, 836, 1
- Herbst, E. 2017, *International Reviews in Physical Chemistry*, 36, 287
- Huang, J., Öberg, K. I., Qi, C., et al. 2017, *The Astrophysical Journal*, 835, 1

- Hudson, R. L., Moore, M. H., Dworkin, J. P., Martin, M. P., & Pozun, Z. D. 2008, *Astrobiology*, 8, 771
- Jørgensen, J. K., van der Wiel, M., Coutens, A., et al. 2016, 595, A117
- Kaifu, N., Ohishi, M., Kawaguchi, K., et al. 2004, *Publications of the Astronomical Society of Japan*, 56, 69
- Kamp, I., Scheepstra, A., Min, M., Klarman, L., & Riviere-Marichalar, P. 2018, *A&A*, 617, A1
- Kral, Q., Wyatt, M., Carswell, R. F., et al. 2016, *Monthly Notices of the Royal Astronomical Society*, 461, 845
- Laas, J. C., Garrod, R. T., Herbst, E., & Widicus Weaver, S. L. 2011, *The Astrophysical Journal*, 728, 71
- Matrà, L., Dent, W. R. F., Wyatt, M. C., et al. 2016, *Monthly Notices of the Royal Astronomical Society*, 464, 1415
- McClure, M., Bailey, J., Beck, T., et al. 2017, *IceAge: Chemical Evolution of Ices during Star Formation*, Tech. rep.
- McClure, M. K., Espaillat, C., Calvet, N., et al. 2015, *The Astrophysical Journal*, 799, 162
- McGuire, B. A., Burkhardt, A. M., Kalenskii, S. V., et al. 2018, *Science*, 359, 202
- McGuire, B. A., Carroll, P. B., Dollhopf, N. M., et al. 2015, *The Astrophysical Journal*, 812, 1
- McGuire, B. A., Shingledecker, C. N., Willis, E. R., et al. 2017, *The Astrophysical Journal Letters*, 851, L46
- Mendoza, E., Lefloch, B., Lopez-Sepulcre, A., et al. 2014, *Monthly Notices of the Royal Astronomical Society*, 445, 151
- Neill, J. L., Muckle, M. T., Zaleski, D. P., et al. 2012, *The Astrophysical Journal*, 755, 153
- Öberg, K. I., Guzmán, V. V., Furuya, K., et al. 2015, *Nature*, 520, 198
- Öberg, K. I., Murray-Clay, R., & Bergin, E. A. 2011, *The Astrophysical Journal*, 743, L16
- Peng, R., Langer, W. D., Velusamy, T., Kuiper, T. B. H., & Levin, S. 1998, *The Astrophysical Journal*, 497, 842
- Pontoppidan, K. M., Salyk, C., Blake, G. A., & Käufel, H. U. 2010, *The Astrophysical Journal*, 722, L173
- Richards, S. N., Moseley, S. H., Stacey, G., et al. 2018, *Journal of Astronomical Instrumentation*, 07, 1840015
- Ros, K., & Johansen, A. 2013, 552, A137

- Schoonenberg, D., & Ormel, C. W. 2017, 602, A21
- Selina, R. J., Murphy, E. J., McKinnon, M., et al. 2018, in Science with a Next Generation Very Large Array, ed. E. Murphy, 15
- Shingledecker, C. N., & Herbst, E. 2018, Physical Chemistry Chemical Physics, 20, 5359
- Trapman, L., Miotello, A., Kama, M., van Dishoeck, E. F., & Bruderer, S. 2017, A&A, 605, A69
- Wakelam, V., Loison, J. C., Herbst, E., et al. 2015, The Astrophysical Journal Supplement Series, 217, 1
- Walsh, C., Millar, T. J., Nomura, H., et al. 2014, 563, A33
- Walsh, C., Loomis, R. A., Öberg, K. I., et al. 2016, The Astrophysical Journal Letters, 823, L10
- Wang, H., Bell, R. C., Iedema, M. J., Tsekouras, A. A., & Cowin, J. P. 2005, The Astrophysical Journal, 620, 1027