

Traces of past activity in the Galactic Centre



Gabriele Ponti

Marie Curie Fellow

Max Planck Institute for Extraterrestrial Physics (Munich)

Morris (UCLA), Terrier, Goldwurm, Clavel, Soldi & Trap (APC Paris)

Cosmic rays in star forming environments - [arXiv:1210.3034](https://arxiv.org/abs/1210.3034)

Was Sgr A* brighter in the past?

$$L_{\text{SgrA}^*} \sim 3 \times 10^{33} - 10^{35} \text{ erg s}^{-1}$$


$$L_{\text{Edd}} = 5 \times 10^{44} \text{ erg s}^{-1}$$

$$M_{\text{SgrA}^*} = 4.4 \times 10^6 M_{\text{Sun}}$$

→ Smallest Eddington ratio BH ($\eta \sim 10^{-9,10}$)

Was Sgr A* brighter in the past?

Map of the Central Molecular Zone (CMZ) in HCN transition




Jones et al. 2012
MOPRA CMZ project

← ~300 pc →



Sgr A* sits on Central Molecular Zone (CMZ) centre

10^7 - $10^8 M_{\text{Sun}}$ of gas in the central 300 pc

- MC tool to trace GC past X-ray activity!
- Correlated variations if clouds along same light front
- Tomography of cold gas distribution!




Clouds to trace Galactic Centre past activity:



→ Link between source incident flux and reflection intensity

$$L_{SgrA^*} \propto \frac{n_H \times A \times I_{FeK}}{d^2}$$

Measure of $d_{\text{line of sight}} \rightarrow L_{\text{source}} \text{ \& } t_{\text{flare}}$



→ Reflection variability modulated by cloud light crossing time
→ Smaller clouds tests faster variability

Related to cloud N_H + los absorption

See e.g. Ryu et al. 2013


Compton scattering depends on source-cloud-observer angle → info on geometry

See e.g. Capelli et al. 2012

FeK & Hard X-ray emission from MC


Which is the origin of the Hard X-ray emission from MC?

GRANAT: Hard X-ray/MC



Sunyaev et al. 1993


ASCA: Fe K α from some MC



Koyama et al. 1996


Are MC reflecting GC radiation?

INTEGRAL: MC - reflection




Revnivtsev et al. 2004

FeK α : constant intensity!



Spectrum:
Sgr B2 consistent with reflection
Luminous event $\rightarrow$ X-ray binary
unlikely

Chandra: Sgr A cont. variability




Muno et al. 2007

But weak detection of variability!!


Alternatives to reflection: Cosmic Rays

Cosmic ray electrons



Contours 850-micron; FeK map (red); 20 cm (green)

Yusef-Zadeh et al. 2002; 2007; 2013



X-ray from MC: Reflection or Cosmic Rays?

⇒ Low energy cosmic ray electrons or protons

Low energy cosmic ray electrons: Yusef-Zadeh et al. 02; 07; 13 Valinia et al. 00;

Low energy cosmic ray protons: Dogiel et al. (2009)

Supernovae ejecta: Bykov 02



⇒ Sgr A* flare $\sim 1.5 \times 10^{39} \text{ erg s}^{-1} \sim 100 \text{ yr ago}$

Internal source

External source: Sgr A*, X-ray binary, Magnetar, Sgr A East


The XMM view of the Galactic Centre

Jones et al. 2012
MOPRA CMZ project



The XMM view of the Galactic Centre


Jones et al. 2012
MOPRA CMZ project



New XMM scan of CMZ (2012)
(continuum)


Red: 0.5-2 keV; Green: 2-4.5 keV; Blue: 4.5-12 keV

Ponti et al. in prep.



The XMM view of the Galactic Centre

Jones et al. 2012
MOPRA CMZ project



New XMM scan of CMZ (2012)
(Fe K emission)

Ponti et al. in prep.

Red: 5-6 keV (continuum); Green: Fe K α emission (reflection); Blue: Fe XXV (hot gas-unresolved sources)


Sgr D

Sgr B

Sgr A

Sgr C


Sgr B2



PRELIMINARY


~300 pc

The X-ray view of Sgr B2




Inui et al. 2009


Terrier et al. 2010




The Chandra-XMM view of Sgr B2



- Luminosity $> 10^{39}$ erg s $^{-1}$ for more than 10 years
- X-ray binary unlikely




Terrier et al. sub.



The XMM view of the Galactic Centre

Jones et al. 2012
MOPRA CMZ project



New XMM scan of CMZ (2012)
(Fe K emission)

Ponti et al. in prep.

Red: 5-6 keV; Green: Fe K α emission; Blue: Fe XXV
(continuum) (reflection) (hot gas-unresolved sources)

Sgr D

Sgr B



Sgr A

Sgr C

PRELIMINARY

~300 pc

Super-luminal motion?



Direction toward Sgr A*


Ponti et al. 2010

Regions causally disconnected!!


→ No propagation of single event → Cosmic ray - internal source excluded!

→ External illumination!

How can a superluminal echo happen?



How can a superluminal echo happen?



If the wall is distant 600000 km from the projector
The variation appears on the screen at $v=2c$!

Because the real “physical” variation happens at the level
of the projector and not of the wall

Possible past activity of Sgr A*...

Sgr A* past activity

$$L_{SgrA^*} \propto \frac{n_H \times A \times I_{FeK}}{d^2}$$


50 km s⁻¹: (No FeK line)

$L_{SgrA^*} < 10^{36} \text{ erg s}^{-1}$ past 60-90 yr

Sgr B2 - G0.11-0.11:

$L_{SgrA^*} \sim 1.4 \times 10^{39} \text{ erg s}^{-1}$ ~100 yr ago



Basic unknown: MC distance!



Another possible Sgr A* light curve


Sgr C:

Line of sight distance through X-ray absorption
(absorption uniformly distributed)



Ryu et al. 2013

3-d distribution of
molecular clouds





$L_{\text{Sgr A}^*} \sim 10^{39}$ erg s⁻¹ for ~400 yr
 $L_{\text{Sgr A}^*} < 10^{35}$ erg s⁻¹ last 100 yr

Another possible Sgr A* light curve

Sgr A:


Line of sight distance through Fe K α line equivalent width



A trend of decreasing luminosity in past 150 yr,
Several flares?


$L_{\text{Sgr A}^*} < 10^{35} \text{ erg s}^{-1}$ last 100 yr

Sgr A's past activity*



A past major outburst of Sgr A?*

Fermi data reveal giant gamma-ray bubbles



Credit: NASA/DOE/Fermi LAT/D. Finkbeiner et al.


APOD 10/11/2010

- Energy release $>10^{55}$ erg
- $\sim$ few 10^6 yr ago

Origin:

- i) a past accretion event onto Sgr A*
- ii) or a nuclear starburst

An AGN torus in our backyard?



adapted from
Molinari et al. 2011

CMZ obscured Sgr A* equatorial view with up to $N_H \sim 10^{24}-10^{25} \text{ cm}^{-2}$

Typical MC core with $v \sim 10^2 \text{ km s}^{-1}$ at 100 pc has $t_{\text{orbit}} = 10^7 \text{ yr} \rightarrow$ AGN torus structure preserved!

To power AGN phase $2 \times 10^3 M_{\text{Sun}}$ accreted $< 0.1 M_{\text{CND}} \rightarrow$ CND can feed Sgr A* again (clumpy CND structure suggest episodic feeding)

Limit cycle of recurrent nuclear activity: Viscous disc evolution $r_{\text{in CND}}$ shrinks $\rightarrow$ Accretion generate shocks/gas compression $\rightarrow$ Trigger star formation $\rightarrow$ Star winds clear gas from GC creating a cavity $\rightarrow$ After 107 yr stellar winds fade $\rightarrow r_{\text{in CND}}$ shrinks **Morris +99**

Summary:

Sgr A*:

At present is extremely under-luminous

Echo of recent (10^2 yr) past weak (10^{39} erg s $^{-1}$) activity from MC of CMZ:

i) Super-luminal FeK variations $\rightarrow$ external illumination $\rightarrow$ **correlated (delayed) cloud variations**

- $\rightarrow$ Tool to determine the GC past (10^3 yr) X-ray activity
- $\rightarrow$ Tool to determine the cloud distribution within the CMZ

Sgr A* outburst:


one long ($\sim 10^2$ yr) highly variable event or several short (< 2 -10 yr) large flares?

Mechanism for such outburst?

(accretion from star winds, asteroid-planet, interstellar medium blob, partial tidal disruption, Sgr A East, gas supply via star tidal interaction)





AGN activity $\sim 6 \times 10^6$ yr ago:

Is the (outer) CMZ a relic AGN torus?



Can irradiation from external sources explain the emission from ALL MC?

Fukuoka et al. 2009



G0.174-0.233 → EW FeK = 950 eV → Reflection

G0.162-0.217 → EW FeK = 360 eV → LECRE

Regions S&N FeK → Cosmic Ray protons!

Other regions variable FeK emission → X-ray binary

Capelli et al. 2011; Tatischeff et al. 2012

Outburst from a magnetar

April 25th Swift detects $L=1.6 \cdot 10^{35} \text{ erg s}^{-1}$ at Sgr A* position
Degenaar et al ATEL5006

April 26th source still at $L=1.7 \cdot 10^{35} \text{ erg s}^{-1}$ at Sgr A* position
Dwelly, GP ATEL5008

SGR-like flare from Sgr A* position (Swift-BAT)
Kennea et al ATEL5009



NuSTAR detects spin period of 3.76 s
Mori et al ATEL5020

Chandra locates the new magnetar at 3" from Sgr A*
Rea et al ATEL5032

Spin down period → magnetic field $B=3 \cdot 10^{14} \text{ G}$
Gotthelf et al ATEL5046

→ New magnetar!

Nature News; Mori et al 2013;
Kennea et al 2013; Rea et al 2013;
Eatough et al. 2013; Bower et al.
2013; Spitler et al. 2013



The Suzaku view of Sgr B2

